

LA-UR-19-30856

Approved for public release; distribution is unlimited.

Title: Convincing Search for Sterile Neutrinos at LANL Coherent
Captain-Mills (CCM) Experiment

Author(s): Van De Water, Richard G.

Intended for: APS Meeting, 2019-04-13 (Denver, Colorado, United States)


Issued: 2019-10-24

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Convincing Search for Sterile Neutrinos at LANL

Coherent Captain-Mills (CCM) Experiment



CAPTAIN = "Cryogenic Apparatus for Precision Tests of Argon Interactions with Neutrinos"

LANL Team

P-25, P-23, P-27, AOT

R.G. Van de Water (PI, Spokesperson), Elena Guardincerri (co-PI), Walter Sondheim, Tyler Thornton, En-Chuan Huang, T.J. Schaub, Mitzi Boswell, Bill Louis, Steve Elliot, Charles Kelsey, Charles Taylor, **Dan Poulson, Bob Macek**

T-2: Daniele S. M. Alves (**co-PI**), Joe Carlson, Rajan Gupta

External Team

Mike Shaevitz (Columbia), H. Ray (U. Florida) Janet Conrad (MIT), **Robert Cooper, (LANL-NMSU, Co-Spokes),** Josh Spitz (U. Mich), R. Van Berg (U. Penn), M. Toups (FNAL), R. Tayloe (IU), and many students + PD's
Darryl Smith (Embry Riddle), Alexis Aguilar-Arevalo (UNAM – Mexico)


WHY NOW?

Short baseline anomalies did not go away,
instead, 2018 MiniBooNE + LSND are
consistent and combined $\sim 6\sigma$

An unambiguous experimental test is
needed demonstrating:

$$\left\{ \begin{array}{l} \nu_{\mu} \rightarrow \nu_s \rightarrow \nu_e \\ \nu_e \rightarrow \nu_s \\ \nu_{\mu} \rightarrow \nu_s \end{array} \right.$$

New MiniBooNE + LSND



WHY LANL?

Coherent CAPTAIN-Mills is the only experiment being proposed that can
test ν_{μ} disappearance with sufficient sensitivity at the LSND mass scale

CCM will be complementary to other neutrino programs around the world

CCM is unique, well-motivated, timely, and fully funded by LANL LDRD funding!



Testing the sterile neutrino hypothesis

Past, current, and future experiments:

Oscillation Mode	Experiment Type	Past/Current Experiments	Signal Significance at LSND Mass Scale	Future Experiments (next 5 years)
$\nu_\mu \rightarrow \nu_e$	Short baseline accelerator	LSND, MiniBooNE, MicroBooNE	6.1σ	SBN@FNAL program, JSNS ²
$\nu_e \rightarrow \nu_s$	Reactor/source	Daya Bay, RENO, Double Chooz	$\sim 2-3 \sigma$	PROSPECT, DANNS, SOLID, BEST, NEOS
$\nu_\mu \rightarrow \nu_s$	Short/Long baseline accelerator	SciBooNE+Mini-BooNE, MINOS+, IceCube	none	CCM

New experiment proposes 100 kg CsI detector at SNS
[arXiv:1901.08094](https://arxiv.org/abs/1901.08094)

Severe Tension Between Appearance & ν_μ Disappearance Experiments in a 3+1 Model



3+1 Models With ν_e Appearance Require Large ν_e & ν_μ Disappearance!

In general, $P(\nu_\mu \rightarrow \nu_e) \sim \frac{1}{4} P(\nu_\mu \rightarrow \nu_x) P(\nu_e \rightarrow \nu_x)$

Assuming that the 3 light neutrinos are mostly active and the N heavy neutrinos are mostly sterile.

More Exotic SBL Possibilities Than 3+N Models (Sterile neutrinos may have other interactions!)

- Sterile Neutrino Decay
- Sterile Neutrinos NSI & New Gauge Bosons
- Altered Dispersion Relations (Resonant Oscillations)
- Pseudo-Dirac Neutrinos
- Light WIMP Production (Light WIMPs can behave like neutrinos)
- Lorentz Violation & CPT Violation
- Mass-Varying Neutrinos
- Neutrino De-Coherence
- etc.

Require
 $\nu_\mu \rightarrow \nu_s$
measurements at
LSND energy to
resolve different
models

Coherent CAPTAIN-Mills experiment

Production mechanism:


charged pions decaying at rest \Rightarrow monoenergetic neutrinos

$$\begin{array}{ccccc} \xleftarrow{\mu^+} & & \pi^+ & & \xrightarrow{\nu_\mu} \\ & & & & E_{\nu_\mu} = 30 \text{ MeV} \end{array}$$


Detection mechanism:

Coherent Elastic Neutrino-Nucleus Scattering
“CEvNS”

present for all active neutrinos



absent for sterile neutrinos





LANSCCE-Lujan Facility a unique place to perform significant and timely test of Sterile Neutrinos

Lujan Target

- Intense protons (100 μA)
- Fast Beam width (250 nsec)
- Compact source

Intense source muon neutrinos: target MCNP simulation flux $4.74 \times 10^5 \text{ nu/cm}^2/\text{s}$ at 20 m




Lujan Experimental Area

- Space for large 10-ton liquid Argon neutrino detector.
- Run detector in multiple locations.
- Room to deploy shielding, large overhead crane, power, etc

Lujan is a Competitive Neutrino Source


Low duty factor critical for background rejection




- Neutrino experiments require high instantaneous power (signal/background)
SNS = 0.029 kJ/nsec; Lujan= 0.028 kJ/nsec
- Plan to run Lujan at ~100 nsec beam width with minimal intensity reduction⁸

Detecting Coherent Neutrinos: Maximizing Scintillation Light Detection!

- 120 R5912 PMT's, wavelength shifting TPB foils



RAT/GEANT Detector Simulation




Simulations predict ~ 0.5 PE/keVnr

- **Liquid Argon scintillates at 128 nm with 40,000 photon/MeV, or 40 photons/keV.**
 - fast 6 nsec and slower ~ 1.6 usec time constants.
 - TPB wavelength shifting coating on PMT's and foils to convert to visible light.
- **Detailed RAT/GEANT4 simulation predicts 10-20 keV detection threshold.**

Expected CAPTAIN-Mills LAr Event Rates (80 kW @ 8 months, 7 tons LAr)



The cross-section is *large*



Large LAr **coherent elastic neutrino-nucleus scattering (CEvNS)** cross sections -> 1000's events!

Reaction	L = 20 m (events/yr)	L = 40 m (events/yr)
Coherent ν_μ (E = 30 MeV)	2709	677
Coherent $\nu_e + \bar{\nu}_\mu$	9482	2370
Charged Current ν_e	257	64
Neutral Current ν_μ	36	18
Neutral Current $\bar{\nu}_\mu$	79	20


- Two oscillation analysis samples with different strategy/backgrounds:
 - PROMPT** with beam (mono-energetic ν_μ) – scattering end point energy 50 keV
 - DELAYED** 4 usec after the beam ($\nu_e + \bar{\nu}_\mu$) - scattering end point energy 148 keV



Signal/Background Strategy

New Reactor-4 result of $7\text{eV}^2 \rightarrow \sim 5\text{m}$ oscillations



- Looking for up to $\sim 10\%$ disappearance over 15-40 m distance
 - $\sim 1,000$ CEvNS events 3 years.
 - Near/far cancellation to reduce systematic errors.
 - Can move detector to multiple positions (sample L/E).
- **Background mitigation crucial, attack with flexible strategy**
 - Fast ($\sim \text{nsec}$) detector and beam 2.9×10^{-6} duty factor (250 nsec)
 - Variable beam width reduced to 30 to 100 nsec for systematic checks.
 - Instrumented and integrated veto.
 - Beam off subtraction (precise, but affect statistics on signal)
 - Particle ID – separate electron and nucleon events $> 10^2$
 - Shielding – can adapt as neutron background measured.



Beam Neutron Backgrounds

- Neutrons from the target, and interactions in the surrounding material.
- No beam off subtraction and veto provides minimal rejection.
- **Prompt Signal:** EJ-301 detectors measured bulk neutrons < 70 MeV. Expect ~100 nsec (200 nsec) neutron free window near (far) position.
- **Delayed Signal:** Low (slow) energy neutrons efficiently rejected with concrete/water shielding.

Neutrino window free of Neutrons: More upstream steel shielding increases window



Building CCM in ~4 months was a Herculean Team Effort!


Students

Alex Diaz (MIT)

Jose Palata (UNAM)

Nick Kamp (UMich)





Complete PDS System Test at LANL with Coherent Captain-Mills (CCM) Detector

- LAr cold test entire SBND PDS system: 96 TPB coated + 24 uncoated PMT's, mounts, cables, feedthrus, HV, electronics, trigger, DAQ, calibration, simulations and data analysis.
- Built detector August-Dec 2018 at LANSCE/Lujan center (100 kW neutron/stopped pion neutrino source)



TPB coated PMTs

Uncoated PMTs

TPB coated reflector foils.
Maximize light output to detect
coherent neutrino-nucleus scattering




- Built detector August-Dec 2018 at Lujan center (100 kW neutron/stopped pion neutrino source)
- **10 tons of LAr filled Dec 10**
- System stable, low noise, long DAQ runs.
- One bad HV channel found (breakdown).
- 20TB of data taken over ~8 beam days

First Results from System Testing: Scope Traces

- Single PhotoElectron ~ 5 mV
- RMS noise < 0.1 mV

Cosmic Ray




Singlet light ~ 6 nsec

Triplet light ~ 1.6 usec

- Electrons have low singlet/triplet ratio
 - Nucleons have high singlet/triplet ratio
- } Pulse shape discrimination
Particle ID


CCM Lujan Run: Dec 12 to 20th

- Beam stable with high uptime and ~ 82 μA current, total $1.5\text{E}20$ POT
 - 3871036 beam triggers with no steel shielding
 - 2482616 beam triggers with steel shielding
- Detector ran stable,
- LAr loss rate of $\sim 1.5''$ day, require 3 top offs during run.
- DAQ performed well, 20 Tbytes of data taken.




Searching for Single PEs in Pre-Beam Data (T. J. Schaub)


- One dead PMT's. Two with bad SPE gains.
- Can see clear single PE peak and noise wall



Clear Single PE peak




Threshold ~0.3 PEs




Beam Event Definition, Global Event View

- DAQ readout window 16 μsec : 10 μsec pre-beam, 4 μsec post beam.
- Find all PMTs for a given time sample that pass a threshold (~ 0.3 PEs)
- An event is consecutive samples with 5 or more PMTs above threshold
- Amplitude and Integral are calculated as the sum amplitudes and integrals of the individual PMTs for the event range



Sanity check: cosmic-ray muons (Dan Paulson)

- Cosmic ray muons reach the Earth surface at a rate $\sim 1/\text{cm}^2/\text{minute}$
- Their flux is proportional to $\cos^2(\theta)$, θ being the zenith angle
- Most of them are Minimum Ionizing Particles and deposit $\sim 2.1 \text{ MeV/cm}$ in LAr
- Muons going through the top and bottom of the tank deposit $> 300 \text{ MeV}$
 - They produce more than $12\text{E}6$ photons in the tank, HUGE signal!
- Based on the shape and size of the inner tank their expected rate is 331 muon/s
- Muons just crossing the top of the tank are expected with a rate of 555 muons/s





With Simple Cuts

- 334 events compatible with muons found in 1.0 s, **equivalent to muons/s**
- Since we are not requiring PMTs from the top region to fire we **expect a rate between 331 and 555 muons/s**
- **The measured rate is consistent with the expected rate** and possibly some shielding resulting from the east side of the building being below grade.


Ar39 Rates and Energy Calibration

- Beta decay endpoint 565 keV, mid energy 219 keV.
- Scan for events in pre-beam region
- Measured rate $\sim 1.4 \times 10^{-4}$ per 150 nsec window, consistent with expectation.




Beam Events with and without Shielding

- Observe beam neutron turn on relative to speed of light particle (~ 100 nsec)
- More shielding decreases neutron rate and increases timing shift
- Gives confidence more steel shielding will increase neutron free region




CCM Measuring Muon-Neutrino Disappearance with Neutral Current Coherent Neutrino Scattering

$\nu_\mu \rightarrow \nu_s$ Disappearance




- **Prompt 30 MeV monoenergetic muon neutrinos:**
 - L/E with no energy smearing.
 - Source position error 8 cm ($\sim 0.4\%$).
- **Delayed neutrinos**
 - High rate signal, high energy (148 keV), different backgrounds.
- **Neutral Current Coherent Ar Scattering:**
 - Flavor independent, all active neutrinos detected, deficit implies oscillation into sterile neutrinos.



- **CEvNS muon disappearance signal would provide smoking gun proof of sterile neutrinos at the LSND+MiniBooNE mass scale.**

CCM Sensitivity to “3+1” sterile neutrino hypothesis (3-year run)



- Can prove/disprove at $\sim 3\sigma$ LSND 3+1 sterile neutrino hypothesis.
- Five year run would approach 5σ !
- If no signal, can rule out world best fit at better than 90%

Summary

- Build CCM detector in 4 months and ran for a week at the end of 2018.
- Demonstrated stable operations, 0.5 PE/keV response with a PMT based LAr detector.
- Will use TOF to isolate mono-energetic neutrinos from neutron backgrounds.
- Calibration and beam running from May-Dec 2019 will establish CEvNS signal.
- Will add more shielding and push for shorter beam width of 100 nsec to improve signal separation.
- Building upgraded detector CCM200 with twice the photocathode coverage and begin muon neutrino disappearance run in 2020.
- Seeking funding for a second CCM detector to improve sterile neutrino and DM search (NSF, DOE Dark Matter FOA, etc)



Backups: The Neutrino Scatters Here!



Coherent elastic neutrino-nucleus scattering (CEvNS)

$$\nu + A \rightarrow \nu + A$$

A neutrino smacks a nucleus via exchange of a Z , and the nucleus recoils as a whole;
coherent up to $E_\nu \sim 50$ MeV



- **Low energy nucleus recoil $E \sim 10$'s keV**
- Well-calculable cross-section in SM:
SM test, probe of neutrino NSI
- Dark matter direct detection background
- Possible applications (reactor monitoring)


$$\frac{d\sigma}{d\Omega} = \frac{G^2}{4\pi^2} k^2 (1 + \cos \theta) \frac{(N - (1 - 4 \sin^2 \theta_W) Z)^2}{4} F^2(Q^2)$$

$$\propto N^2$$


Coherent Neutrinos have been Recently Observed at SNS

First light at the SNS with 14.6-kg CsI[Na] detector

Charge



Time



Observation of coherent elastic neutrino-nucleus scattering

D. Akimov^{1,2}, J. B. Albert³, P. An⁴, C. Awe^{4,5}, P. S. Barbeau^{4,5}, B. Becker⁶, V. Belov^{1,2}, A. Brown^{4,7}, A. Bolozdy...

+ See all authors and affiliations

Science 03 Aug 2017;
eaao0990
DOI: 10.1126/science.aao0990





Peer Reviewed
← see details



D. Akimov et al., *Science*, 2017


<http://science.sciencemag.org/content/early/2017/08/02/science.aao0990>

MiniBooNE New Oscillation Results



- MiniBooNE is consistent with LSND excess, and combined is 6 σ

Integrated and Active Veto Regions for Background Rejections




- 7 tons LAr Fiducial volume, 3 tons LAr Veto (2-3 radiation lengths).
- Active Veto region crucial to rejecting cosmic rays and other external backgrounds.
 - 1 MeV gamma: 2 rad lengths: 100 keV x-ray: 10 rad lengths: 1 MeV neutron: 1 scatter length

Coherent Neutrino-Nucleus Scattering Energy Spectrum


Prompt Neutrinos $E_{\text{muon}} = 30 \text{ MeV}$

Prompt ν_μ Disappearance ($\sin^2(2\theta) = 0.1$, $L=18 \text{ m}$)




Delayed Neutrinos

Delayed ν_e Disappearance ($\sin^2(2\theta) = 0.1$, $L=18 \text{ m}$)



Delayed $\bar{\nu}_\mu$ Disappearance ($\sin^2(2\theta) = 0.1$, $L=18 \text{ m}$)




Random Backgrounds for CEvNS


- **Cosmogenic:** about ~ 200 Hz. Beam duty factor + overburden + veto, will reduce to 10 (130) events/year.
- **U/Th:** Dust and PMT glass contains U/Th \sim MeV beta/gamma. Expect ~ 10 Hz per PMT. Analysis tricks such as fiducial cuts, or large charge in a single PMT cuts will reduce significantly. Will need to pay attention to cleanliness during detector construction. Expected background small.
- **Beam off subtraction will measure these backgrounds extremely well.**
- 295 nsec beam good, but some running with shorter beam time of 30 or 100 nsec would provide systematic check on background estimates.

Random Backgrounds for CEvNS

- ^{39}Ar : 565 keV endpoint β emitter: In CCM detector, estimate 7.5 kHz rate.
 - Beam rejection factor $\sim 10^5$
 - energy cut ($\sim 10^1$), pulse height discrimination (10^1 to 10^5)
 - Measured precisely by beam off running.
 - Subtraction increases signal statistical error from 2.7% to 3.5% (prompt), minimal effect on delayed signal.



Particle Separation with Pulse Height Analysis



Detailed Target MCNP Simulations (Charles Kelsey P-27)

Nuclear Instruments and Methods in Physics Research A 594 (2008) 373–381
Nuclear Instruments and Methods in Physics Research A 632 (2011) 101–108








Fig. 1. Elevation view of the Lujan Center's TMRS geometry used in our calculations. The main components are labeled: split tungsten target (W), beryllium reflector (Be), lead reflector–shield (Pb), and the steel reflector–shield (Fe).



- Simulations has confirmed hand calculated flux of $\sim 4.74 \times 10^5$ nu/cm²/s at 20 m
- Horizontal extend of neutrino production at the source of 8 cm (1 sigma position error)
- Simulated neutron backgrounds




Initial Neutron Rates and Spectrum (TOF) Measurements with EJ-301 Neutron Detectors



Neutron Reduction with 26" Steel

Neutron time of flight over 10.8 m



**Leading edge of neutrons ~76 MeV
which is 168 nsec delay over 20m**

Timing Delays Measured (Scope): Working out a few missing pieces – dealing with legacy knowledge/equipment

Future upgrades to improved timing delay stability and simplicity, i.e. faster coils, direct cable from coil to detector digitizers, etc

20 m n-TOF table

	c	67 ns
800 MeV		79 ns
500 MeV		88 ns
200 MeV		118 ns
100 MeV		156 ns
50 MeV		213 ns
20 MeV		328 ns

